

TO STUDY THE COMPLICATIONS IN DENGUE LIKE ILLNESSES

Girdhari lal Sharma¹, Rakesh Bilwal²

^{1,2}Assistant Professor, Department of Pediatrics, National Institute of Medical Sciences and Research Jaipur, Rajasthan, India

Article Info: Received 18 June 2020; Accepted 09 July 2020

DOI: <https://doi.org/10.32553/ijmbs.v4i7.1284>

Corresponding author: Rakesh Bilwal

Conflict of interest: No conflict of interest.

Abstract

Background: Dengue has a wide spectrum of clinical presentations, often with unpredictable clinical evolution and outcome. While most patients recover following a self-limiting nonsevere clinical course, small proportion progress to severe disease

Methods: The hospital based study was conducted on patients presenting to tertiary care teaching hospital, who fulfilled inclusion and exclusion criteria were enrolled for the study. A comprehensive history taking, physical examination, and lab investigations were carried out and data were collected in pre-designed proforma.

Results: According to complications, in dengue positive cases, bleeding was the most common complication seen in 48.00% cases while shock, myocarditis, and convulsion were present in 31.00%, 2.00% and 1.00% of cases respectively. Among the dengue negative cases, 60.00% had no complication while shock and bleeding were present in 28% and 12.00% cases respectively.

Conclusion: It concluded that shock was most common complication.

Keywords: Hematocrit, Dengue, Dengue Like Illness

Introduction

Infection with the Dengue virus (DENV) is increasingly recognized as an important arthropod-borne viral infection infecting about 2.5 billion people worldwide; being endemic to over a 100 countries with over 975 million belonging to tropical and sub-tropical countries in Southeast Asia, the Pacific and the America. Dengue virus is a single stranded RNA virus which was first isolated from Japan in 1942 by Hotta; while belonging to Flaviviridae family it is transmitted to humans by infective female of Aedes genus, principally by Aedes aegypti, Aedes albopictus mosquito, Aedes polynesiensis and several species of the Aedes scutellaris complex. There are 4 antigenically distinct serotypes, i.e. DENV-1, DENV-2, DENV-3 and DENV-4, which are evolved from a common ancestor.¹

Dengue is the most important arthropod-borne viral infection of humans. Each year there are about 50 million dengue infections and about 500,000 individuals are hospitalized with DHF, mainly in Southeast Asia. Dengue has been an urban disease but now has spread to rural areas of India as well²

Dengue has a wide spectrum of clinical presentations, often with unpredictable clinical evolution and outcome². While most patients recover following a self-limiting nonsevere clinical course, small proportion progress to severe disease. Early recognition of dengue is challenging because the initial symptoms are often non-specific.

Viremia may be below detectable levels and serological tests confirm dengue late in the course of illness³. The key to a good clinical outcome is understanding and being alert to the clinical problems that arise during the different phases of the disease, leading to a rational approach in case management.

For a disease that is complex in its manifestations, management is relatively simple, inexpensive and very effective in saving lives, so long as correct and timely interventions are instituted. Hence it is very important to be able to predict which patients are more likely to land up in complications. Such early prediction of adverse outcome and early detection of danger signs can help in reducing mortality and morbidity. Predictors will also help in monitoring the therapy and the course of illness in dengue-like illnesses.⁴

Since there is a paucity of literature on this, and there exist many gaps in the current understanding of such predictive factors, this study is being planned to assess history, examination, and investigation based predictive factors for complications of dengue-like illnesses.

Material and Methods

Study Design: Hospital-based cross-sectional study.

Sample size- 200 patients fulfilling the following inclusion criteria:

All the patients admitted to Paediatric ward with-

1. Age 3 months to 14 years.

2. A history of fever as given by the patient or parents or a documented fever > 38-degree Celsius in the first 24 hours after admission.

3. Thrombocytopenia of < 100,000 /cubic mm as documented in the automated counter within 24 hours of admission.

4. Informed consent obtained from the guardians or relatives.

Exclusion criteria

1. Patients with a proven bacterial infection as the cause for thrombocytopenia and fever.

2. Patients with a pyogenic focus of infection.

3. Discharge against medical advice or referral to higher center before ascertaining the final outcome.

The study was commenced after obtaining clearance from the institutional ethical committee. Written consent was taken from the parents and those who were not willing excluded from the study. Patients presenting to paediatric hospital, who fulfilled inclusion and exclusion criteria were enrolled for the study. A comprehensive history taking, physical examination, and lab investigations were carried out and data were collected in pre-designed proforma.

OBSERVATIONS

Table 1: Socio-demographic variable

Variable	Dengue		p-value
	Positive	Negative	
Age in Years (Mean±SD)	10.12±3.16	10.22±3.19	0.123
Male : Female	61 : 39	64 : 36	0.256

Mean age was 10.12±3.16 year in dengue negative cases and 10.22±3.19 year in dengue positive cases. This difference was found statistically insignificant ($p>0.05$). In the present study, male patients outnumber than the female with a male to female ratio of 1.87:1. This difference was found statistically insignificant ($p>0.05$).

Table 2: Distribution of Cases According to Complication

Complication	Dengue			
	Negative		Positive	
	No.	%	No.	%
Bleeding	12	12.00	48	42.00
Myocarditis	0	-	2	2.00
Shock	28	28.0	31	31.00
Convulsion	0	-	1	1.00
No Complication	60	60.00	18	12.00
Total	100	100.00	100	100.00
P	<0.001			

According to complications, in dengue positive cases, bleeding was the most common complication seen in 48.00% cases while shock, myocarditis, and convulsion were present in 31.00%, 2.00% and 1.00% of cases

respectively. Among the dengue negative cases, 60.00% had no complication while shock and bleeding were present in 28% and 12.00% cases respectively.

On applying the chi-square test, the difference was found statistically highly significant ($p<0.001$).

Discussion

Dengue is a major international health concern that is prevalent in tropical and sub-tropical countries. Since the first confirmed case of dengue in India, during the 1940s, intermittent reports from Delhi, Ludhiana, Mangalore, Vellore and from other states have been published. The diagnosis is by clinical profile but they can present with varied manifestation⁵⁻⁶.

There is a steady increase in the outbreak of dengue fever over the years and so among children. This is due to the rapid urbanization with unplanned construction activities and poor sanitation facilities contributing fertile breeding grounds for mosquitoes. Due to an increase in the awareness among health care professionals following the initial epidemic and the availability of diagnostic tests have contributed to the increased diagnosis⁷.

A outbreak of dengue fever during pre-monsoon and monsoon season reported due to stagnation of water after a bouts of rainfall which facilitate vector breeding. This highlight the preventive measures against dengue fever should be taken during water stagnation periods after the initial bouts of rainfall and at the end of monsoon.

Mean age was 10.12±3.16 year in dengue negative cases and 10.22±3.19 year in dengue positive cases. This difference was found statistically insignificant ($p>0.05$). This may be due to out-door activities of these children, where chances of getting bitten by mosquitoes are more. Similar finding was observed in other studies⁸⁻¹⁰.

In the present study, male patients outnumber than the female with a male to female ratio of 1.87:1. This difference was found statistically insignificant ($p>0.05$).

Boys were slightly more affected then girls were also observed by Selvan et al¹⁰ and Sahana et al⁹ and similar pattern was seen in the retrospective analysis of the 2006 North Indian Dengue outbreak¹⁴. This may be due to out-door activities of these children, where chances of getting bitten by mosquitoes are more.

According to complications, in dengue positive cases, bleeding was the most common complication seen in 48.00% cases while shock, myocarditis, and convulsion were present in 31.00%, 2.00% and 1.00% of cases respectively. Among the dengue negative cases, 60.00% had no complication while shock and bleeding were present in 28% and 12.00% cases respectively.

Raj et al¹¹ observed that shock was the most common and difficult to treat complication despite appropriate fluid management in accordance with WHO regimen. 20(10.2%) children had shock, of which 7 were refractory to fluid therapy and blood products (whole blood, packed cell volume, FFP) given as indicated.

Conclusion

It concluded that shock was most common complication.

References

1. Guzman MG, Halstead SB, Artsob H, Buchy P, Farrar J, Gubler DJ, Hunsperger E, Kroeger A, Margolis HS, Martinez E, et al. Dengue: a continuing global threat. *Nat Rev Microbiol.* 2010;8:S7–16. doi: 10.1038/nrmicro2460.
2. Halstead SB. More dengue, more questions. *Emerg Infect Dis* 2005; 11(5):740-41.
3. Ramos MM, Tomashek KM, Arguello DF, Luxemburger C, Quinones L et al. Early clinical features of dengue infection in Puerto Rico. *Transactions of the Royal Society of Trop Med Hygiene* 2009; 103(9):878-84.
4. Chaudhary R, Khetan D, Sinha S, Sinha P, Sonker A et al. Transfusion support to dengue patients in a hospital base blood transfusion service in North India. *Transfusion and Apheresis Sci* 2006;3 5(3);239-44.
5. Padibidri VS, Adhikari P, Thakare JP, Ilkal MA, Joshi GD, Pereira P, et al. The 1993 epidemic of dengue fever in Mangalore, Karnataka State, India. *Southeast Asian J Trop Med Public Health.* 1995;26:699-704.
6. Cherian T, Ponnuraj E, Kuruvilla T, Kirubakaran C, John TJ, Raghupathy P. An epidemic of dengue hemorrhagic fever and dengue shock syndrome in and around Vellore. *Indian J Med Res.* 1994;100:51-6.
7. Gubler DJ. Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev.* 1998;11:480-96.
8. Mittal H, Faridi MM, Arora SK, Patil R. Clinicohematological profile and platelet trends in children with dengue during 2010 epidemic in north India. *Indian J Pediatr.* 2012;79:467-71.
9. Sahana KS, Sujatha R. Clinical profile of dengue among children according to revised WHO classification: analysis of a 2012 outbreak from Southern India. *Indian J Pediatr* 2015;82:109-13.
10. Selvan T, Nagaraj MV, Saravanan P, Somashekar. A study of clinical profile of dengue fever in children. *Int J Contemp Pediatr* 2017;4:534-7.
11. Raj AS, Munshi S, Shah BH. A study on clinical presentation of dengue fever in Children. *IJMSR,* 2016; 2272-8.