

Comparative Study of Different Doses of Dexmedetomidine As An Adjuvant To Intrathecal Hyperbaric Bupivacaine In Lower Limb Orthopaedic Surgeries

Dr Rekha Verma¹, Dr Ranjeet²

¹MBBS, MD (Anaesthesia)

²MBBS, MS (ENT)

Article Info: Received 05 August 2022; Accepted 07 September 2022

doi: <https://doi.org/10.32553/ijmbs.v6i9.2603>

Corresponding author: Dr Rekha Verma

Conflict of interest: No conflict of interest.

Abstract

Background: This study is aimed to assess the effect of Intrathecal administration of different doses of Dexmedetomidine with hyperbaric Bupivacaine.

Methods: A prospective randomized double blind study was conducted with 150 consenting patients of ASA grade I and II, scheduled for lower limb Orthopaedic surgeries. Using the lottery method, the patients were randomly allotted into 3 groups, 50 patients in each group. Group A, Group B & Group C.

Results: Post-operative VAS and total analgesic requirement in 24 hours were minimal in group C as compare to B group. All the patients achieved Bromage scale 3 motor blocks and there was dose dependent prolongation of motor block in B and C groups. Similarly regression of motor block to Bromage 0 was significantly prolonged in group C than B and A group. Complete recovery of sensory and motor functions was observed in all the patients.

Conclusion: Supplementation of spinal Bupivacaine with Dexmedetomidine significantly prolonged both sensory and motor block compared with intrathecal Bupivacaine alone.

Keywords: Dexmedetomidine, Bupivacaine, Intrathecal

Introduction

Pain is an unpleasant feeling often caused by intense or damaging stimuli. It has been defined by the International Association for the Study of Pain (IASP) as “An unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage”.¹⁻³

Dexmedetomidine is highly selective α_2 adrenergic agonist. Dexmedetomidine has been used as intrathecally as an adjuvant and no neurological side-effect is reported in humans. It also provides stable hemodynamic condition, good quality of intra-operative and prolonged post-operative analgesia with minimal side effects. Intrathecal α_2 receptor agonists are found

to have antinociceptive action for both somatic and visceral pain.⁴

Material and Method

Type of study: A prospective randomized double blind study

Inclusion criteria:

Patients of ASA grade I and II, scheduled for lower limb Orthopaedic surgeries.

Exclusion criteria:

1. Patients with hypotension, coagulation defects, spine abnormalities, heart block, arrhythmias etc.
2. Body weight ≥ 120 kg and height ≤ 150 cm.

3. Patients on calcium channel blockers, adrenergic receptor blockers, and ACE inhibitors.

A prospective randomized double blind study was conducted with 150 consenting patients of ASA grade I and II, scheduled for lower limb orthopaedic surgeries. Using the lottery method, the patients were randomly allotted into 3 groups, 50 patients in each group. Group A, Group B &

Group C. The surgeon, patient and the observing anaesthesiologist were blinded to the patient group.

Data analysis

All data were analyzed by Epi-info software. Student t test and ANOVA test for parametric data. Chi square test for non-parametric data.

Result

Table 1: Socio-demographic variable

Variable	Group-A	Group-B	Group-C	P-value
Age in Yrs	34.26±8.02	36.25±7.89	34.26±8.05	>0.05
Male : Female	37:13	36:14	33:17	>0.05
ASA (I:II)	47:3	46:4	45:5	>0.05

All three group were comparable.

Table 2: Out come

Variable	Group-A	Group-B	Group-C	P-value
Time of onset of sensory block	7.23±1.59	8.23±2.64	8.15±2.39	>0.05
Time of onset of motor block	9.36 ±3.39	9.15±2.59	9.11±2.29	>0.05
Duration of sensory block	101.09 ±16.05	115.05±19.23	146.23±20.01	0.01
Duration of motor block	162.02 ±17.56	197.26±25.06	270.23±23.02	0.01

Post-operative VAS and total analgesic requirement in 24 hours were minimal in group C as compare to B group. All the patients achieved Bromage scale 3 motor blocks and there was dose dependent prolongation of motor block in B and C groups. Similarly regression of motor block to Bromage 0 was significantly prolonged in group C than B and A group. Complete recovery of sensory and motor functions was observed in all the patients.

Discussion

Van Tuijl I⁵ added various doses of Clonidine (0, 15 or 30 µg) to 5 mg hyperbaric Bupivacaine and evaluated their effect on the duration of the motor block, analgesic quality and ability to void. They opined that addition of 15 and 30 µg of Clonidine increased the motor block duration by

25 and 34 min, respectively and also resulted in better analgesic quality.

Hutschala D, Mascher H et al⁶ added Clonidine to Bupivacaine and found that it enhances and prolongs analgesia after brachial plexus block via a local mechanism in healthy volunteers.

Niemi L et al⁷ studied effects of intrathecal Clonidine on duration of Bupivacaine spinal anesthesia, hemodynamics, and postoperative analgesia in patients undergoing knee arthroscopy and found that intrathecal Clonidine significantly prolongs the anesthetic and analgesic effects of Bupivacaine. Kalso A(4)reported that as compared to Clonidine, the affinity of DXM to [alpha] 2 receptors is ten times greater. Results of our study showed that

addition of Dexmedetomidine to Bupivacaine although delays onset but, significantly prolongs the duration of sensory and motor block.

Mahmoud M. Al-Mustafa et al⁸ added Dexmedetomidine to spinal Bupivacaine for urological procedures. He compared 5mcg (Group D 5) and 10 mcg (Group D 10) of Dexmedetomidine added to 12.5 mg Bupivacaine to Bupivacaine 12.5 mg with normal saline (Control group). The author found that the mean time of sensory block to reach T10 dermatome was 4.7 ± 2.0 minute in D10 group, 6.3 ± 2.7 minute in D5 group and 9.5 ± 3.0 minute in control group. The mean time to reach bromage 3 scales was 10.4 ± 3.4 minute in D10 group, 13.0 ± 3.4 minute in D5 group and 18.0 ± 3.3 minute in control group. Regression time to reach S1 dermatome was 338.9 ± 44.8 minute in D10 group, 277.1 ± 33.2 minute in D5 group and 165.5 ± 32.9 minute in control group. Time to reach bromage 0 was 302.9 ± 36.7 minute in D10 group, 246.4 ± 24.7 minute in D5 group and 140.1 ± 32.3 minute in control group. They found that Dexmedetomidine has dose dependent effect on onset and regression of sensory and motor block.

Conclusion

Supplementation of spinal Bupivacaine with Dexmedetomidine significantly prolonged both sensory and motor block compared with intrathecal Bupivacaine alone.

References

1. James Leonard Corning and the early history of spinal puncture *Neurology* April 1987 vol. 37 no. 4 672 Marx, GF (1994). "The first spinal anesthesia. Who deserves the laurels?" *Regional Anesthesia*. 19 (6): 429–30.
2. Wang, Josef K. M.D.; Nauss, Lee A. M.D.; Thomas, Juergen E. M.D. Pain Relief by Intrathecally Applied Morphine in Man. *Anesthesiology*(1979);50; 149-151.
3. Kashif M. Madani, Mohit Somani, Khyyam Moin, Sudhir Sachdev, Durga Jethava, Vijay Mathur Comparative study of different doses of Dexmedetomidine in spinal anaesthesia in lower limb orthopaedic procedures. *J of Evolution of Med and Dent Sci*, 2014;3(62):13723-26.
4. Van Tuijl I, Giezeman MJ, Braithwaite SA, Hennis PJ, Kalkman CJ and van Klei WA (2008). Intrathecal low-dose hyperbaric Bupivacaine-Clonidine combination in outpatient knee arthroscopy: a randomized controlled trial. Publication date and source *Acta Anaesthesiol Scand* 52 (3) 343-349.
5. Hutschala D, Mascher H, Schmetterer L, Klimscha W, Fleck T, Eichler HG, Tschernko EM (2004). Clonidine added to Bupivacaine enhances and prolongs analgesia after brachial plexus block via a local mechanism in healthy volunteers. *European Journal of Anaesthesiology* 21 (3) 198-204.
6. Niemi L (1994). Effects of intrathecal Clonidine on duration of Bupivacaine spinal anesthesia, hemodynamics, and postoperative analgesia in patients undergoing knee arthroscopy. *Acta Anaesthesiol Scand* 38724-728.
7. Kalso E, Poyhia R and Rosemberg P (1991). Spinal antinociceptive by Dexmedetomidine a highly selective 2-adrenergic agonist. *Pharmacology Toxicology* 68 140-143.